

Small-angle scattering on a system of magnetic and electric charges

M. Penttinen[◇]

Institut für Theoretische Physik II

Ruhr-Universität Bochum

44780 Bochum, Germany

February 1, 2008

Abstract

The generalization of the conformal scattering method for small-angle scattering processes involving magnetic monopoles and ordinary charges is constructed. Using this generalization we show that introducing of magnetic charges corresponds to analytical continuation of the eikonal amplitude in the complex charge plane (the imaginary part is proportional to the magnetic charge). We calculate explicitly the eikonal amplitude for scattering on a dyon and two monopoles in terms of confluent hypergeometric functions. The singularities of the corresponding amplitudes (focal points) are studied in details.

[◇]DAAD fellow, on leave of absence from St. Petersburg University, St. Petersburg, Russia

1 Introduction

It is known [1], that if the potential energy satisfies the Laplace equation, the small-angle scattering is closely related to conformal mapping on a complex plane. This makes it possible to introduce complex variables in the impact-parameter plane (b) and in the momentum-transfer plane (p), which leads to a remarkable simplification of the calculations. Besides the simplification for harmonic scatterers, it is possible to formulate certain rigorous theorems that permit to estimate the classical scattering at small and large momentum transfers, as well as to elucidate the nature of the singularities in the differential effective cross section.

In the present paper we extend the method of conformal scattering [1, 2] to the case of scattering processes involving magnetic monopoles (dyons) and ordinary charges.

The hypothesis of a new particle, the magnetic monopole, has been introduced by Dirac [3] in 1931. Such a particle, for which there is no experimental evidence would be a source of Coulomb-like magnetic field. Dirac argued that in order to fit into conventional quantum mechanics, the new particle must satisfy a quantization condition:

$$eg = n/2 \tag{1}$$

where g is the charge of the monopole, e is the electric charge of any other particle, and n is some integer. Due to the quantization condition (1) a perturbative treatment of the scattering processes involving magnetic monopoles and usual charges is doubtful if at all possible. In this situation one looks for methods of calculating the scattering amplitude which although approximate, are nonperturbative.

The eikonal approximation [4] for small angle scattering of fast particles not only makes it possible to express the scattering amplitude as an integral, but

also preserves many important properties of the exact solution. For example, it preserves the unitarity relation, which is violated by e.g. the Born approximation. In limiting cases the eikonal approximation reduces to the Born or classical approximations for small-angle scattering.

The small-angle scattering problem simplifies significantly in the case of harmonic scatterers (i.e. when the potential energy satisfies the Laplace equation) due to the analyticity of complex impact-parameter plane mapping to momentum-transfer plane [1, 2]. After the complex variables, namely the impact parameter b and the momentum transfer p , are introduced the integrations in the eikonal formula can be separated and therefore the scattering amplitude factorizes, i.e., it can be expressed as a sum each term of which is the product of two functions depending on p and p^* , respectively. Thus the scattering amplitude can be expressed through contour integrals, what is very convenient for both general studies and numerical calculations.

The outline of the paper is as follows. In section 2 we formulate the method of conformal scattering for processes involving magnetic and electric charges. We show that the description of small-angle scattering processes on a system of electric and magnetic charges can be simplified by introducing the complex charge planes (the electric charge corresponds to the real part and the magnetic charge to the imaginary part). This implies that the amplitudes for scattering on various configurations of electric and magnetic charges are related by analytical continuation of the amplitudes in the complex charge plane. In section 3 the quantum mechanical generalization of the method of conformal scattering is considered. Using this method we examine in details scattering on a dyon and two monopoles located at different points. For the first time we obtain explicit expressions for the eikonal amplitude for all these problems in terms of confluent hypergeometric

functions and Bessel functions of various types for all these problems.

In section 4 we investigate the focal points, where the classical scattering amplitude becomes infinite, which play an important role in the general theory of diffraction and in catastrophe theory.

2 Classical conformal scattering

To begin with, let us consider according to [1] the classical small-angles scattering of particles by nonspherically symmetrical scatterer with potential energy $U(x, y, z)$ in the presence of an external magnetic field with potential $A_i(x, y, z)$. It can be regarded as a one-to-one single-valued mapping of the impact-parameter plane $b(x, y)$ into the transverse momentum-transfer plane $p_\perp(p_x, p_y)$, given by

$$p_\perp = -p_0^{-1} \nabla V(x, y),$$

$$V(x, y) = \int_{-\infty}^{\infty} U(x, y, z) dz - \frac{ep_0}{c} \int_{-\infty}^{\infty} A_z(x, y, z) dz,$$

where e and p_0 are the charge and the momentum of the incident particle, and $A_x, A_y \rightarrow 0$ as $z \rightarrow \infty$ has been assumed. The differential effective small-angle cross section is defined as the Jacobian of the transformation $b \rightarrow p_\perp$

$$\sigma(p_x, p_y) = p_0^2 \sum \left| \frac{\partial(x, y)}{\partial(p_x, p_y)} \right|,$$

where the summation is over all the values of the multiply valued function $b(p_\perp)$ (it can be “zero-valued” for certain $p_\perp$). In other words, the effective cross section is defined as the ratio of the area elements in the plane b and the corresponding area elements in the plane $p_\perp$.

In the general case, a small circular area in the b plane is mapped onto an elliptic area in the $p_\perp$ plane. The line along which the minor semi-axis of this

ellipses vanishes is called the rainbow line (then the major semi-axes are the tangents of the rainbow line). The mapping of the rainbow lines on the impact-parameter plane is defined by the equation

$$\partial(p_x, p_y)/\partial(x, y) = 0. \quad (2)$$

For a spherically symmetrical potential the scattering is axisymmetric and the rainbow lines are circles in the impact-parameter plane and in the momentum-transfer plane with the center at the origin. If the potential $V(x, y)$ is a harmonic function, i.e., it satisfies the two-dimensional Laplace equation $\Delta V(x, y) = 0$ then rainbow lines degenerate into points (the so called focal points), since equation (2) being supplemented by $\Delta V(x, y) = 0$ takes the form:

$$\left(\frac{\partial^2 V}{\partial x^2}\right)^2 + \left(\frac{\partial^2 V}{\partial x \partial y}\right)^2 = 0.$$

Hence instead of one condition that defines the rainbow line on a plane, we obtain thus two conditions that define in the general case only points on the impact - parameter plane and on the momentum - transfer plane (focal points).

It is convinient to introduce complex impact parameter and complex transverse momentum

$$b = x + iy, \quad p = p_x + ip_y$$

as well as a complex potential analytical function, such that $V(x, y) = \text{Re } V(b)$. We then obtain the following formulas for the momentum transfer, the focal point, and the effective cross section

$$p^* = -p_0^{-1} \frac{dV}{db}, \quad \frac{d^2 V}{db db^*} = 0, \quad \sigma = p_0^2 \sum \left| \frac{db}{dp^*} \right|^2, \quad (3)$$

where the summation is over all the values of the multiply valued function db/dp^* .

Let us consider a system of N point-like electric charges q_i situated at points with coordinates $\vec{w}_i$ ($i = 1, \dots, N$) and of M point like magnetic charges g_i at the

points $\vec{z}_i$ ($i = 1, \dots, N$).¹ For such system:

$$U(x, y, z) = \sum_{i=1}^N \frac{eq_i}{|\vec{r} - \vec{w}_i|}, \quad (4)$$

$$\vec{A}(x, y, z) = \sum_{j=1}^M g_j \cdot \frac{[(\vec{n}_j \times (\vec{r} - \vec{z}_j)) \cdot (\vec{n} \cdot (\vec{r} - \vec{z}_j))]}{|\vec{r} - \vec{z}_j|((\vec{r} - \vec{z}_j)^2 - (\vec{n} \cdot (\vec{r} - \vec{z}_j))^2)}. \quad (5)$$

We use a form of the vector potential of magnetic monopole suggested by Dirac [3]

$$\vec{A}(z) = g \cdot \frac{[\vec{n} \times \vec{r}] \cdot (\vec{n} \cdot \vec{r})}{|\vec{r}|(r^2 - (\vec{n} \cdot \vec{r})^2)}. \quad (6)$$

Here $\vec{n}$ is a unit vector in the direction of the “Dirac string” being a line of singularity of the potential (6). The two-dimensional potential corresponding to (4) and (5) has the form

$$V(x, y) = V_{el}(x, y) + V_{mag}(x, y) \quad (7)$$

where

$$V_{el}(x, y) = \int_{-\infty}^{\infty} U(x, y, z) dz = \sum_{i=1}^N q_i \ln(|\vec{b} - \vec{b}_i|^2) \quad (8)$$

with $b_i \equiv (w_{x_i}, w_{y_i})$ being position of the electric charges in a impact parameter plane, $\vec{b} = (x, y)$, and

$$\begin{aligned} V_{mag}(x, y) &= -\frac{ep_0}{c} \int_{-\infty}^{\infty} A_z(x, y, z) dz = \\ &= -\frac{ep_0}{c} \sum_{j=1}^M g_j \cdot \arctan \left(\frac{(\vec{b} - \vec{b}_j) \cdot \vec{n}_j}{[(\vec{b} - \vec{b}_j) \times \vec{n}_j]_z} \right). \end{aligned} \quad (9)$$

Here $\vec{b}_j = (z_{x_j}, z_{y_j})$ are the positions of the magnetic charges, and we choose $n_{jz} = 0$ to satisfy a conditions $A_x, A_y \rightarrow 0$ as $z \rightarrow \infty$. It is easy to check that

¹The case when $\vec{z}_i = \vec{w}_j$ for some i and j corresponds to a dyon with charges (q_i, g_i) at the point $\vec{z}_i = \vec{w}_j$

$V(x, y)$ is a harmonic function. Introducing a complex impact-parameter plane and a complex potential plane we see that

$$\begin{aligned} V_{el}(x, y) &= \operatorname{Re} \sum_{i=1}^N q_i \ln(b - b_i) \\ V_{mag}(x, y) &= -\frac{qp_0}{c} \operatorname{Im} \sum_{i=1}^N eg_i \ln[(b - b_i) \cdot e^{i\theta_j}] \end{aligned}$$

with $\theta_j = \arctan(n_{x_j}/n_{y_j})$. The effective complex potential for a system of N electric and M magnetic charges has the form:

$$V(b) = V_{el}(b) + V_{mag}(b) = \sum_{k=1}^{M+N} Q_k \ln[(b - b_k) \cdot e^{i\theta_k}], \quad (10)$$

where $Q_k = eq_k - \frac{iep_0}{c}g_k$ are complex charges with real part equal to the electric charge q_k at point b_k , and with imaginary charge part proportional to the magnetic charge at the same point. We see that introducing magnetic charges into a system corresponds to continuation of the charges into a complex plane, this fact reduces the problem of scattering on a system involving magnetic charges to the corresponding problem with only electric charges by analytical continuation in the complex charge plane. It is also worth noting that the imaginary part of the complex charges is proportional to the momentum of the incident particle.

The general formulas (3) and (10) being applied to the case of scattering on a point-like dyon with charges q and g give a Rutherford formula

$$\sigma = \frac{1}{|p|^4} \cdot \left\{ \left(\frac{2p_0}{c} eg \right)^2 + (eq)^2 \right\}. \quad (11)$$

As a next example let us consider the problem of scattering by two dyons with complex charges $Q_{1,2} = eq_{1,2} + ieg_{1,2}p_0/c$ at the points $b_1 = R$, $b_2 = -R$. In this case equations (3) and (10) give

$$p^* = \frac{1}{p_0} \cdot \left(\frac{Q_1}{b - R} + \frac{Q_2}{b + R} \right). \quad (12)$$

The equation for the focal points reads:

$$\frac{Q_1}{(b_f - R)^2} + \frac{Q_2}{(b_f + R)^2} = 0 \quad (13)$$

and has the following solutions:

$$b_f^{1,2} = -R \cdot \frac{(\sqrt{Q_1} \pm i\sqrt{Q_2})^2}{Q_1 + Q_2}. \quad (14)$$

The corresponding values of the momentum transfer, where the cross section is infinitely large, are

$$p_{1,2}^f = -\frac{1}{4Rp_0} \cdot \frac{(Q_1 + Q_2)^2}{(\sqrt{Q_1} \pm i\sqrt{Q_2})^2} \quad (15)$$

and the cross section can be written in the form:

$$\sigma = \frac{|Q_1 + Q_2|^2}{2|p|^4} \cdot \left[1 + \frac{1}{4} \left| 2 + \frac{p/p_1^f - 1}{p/p_2^f - 1} + \frac{p/p_2^f - 1}{p/p_1^f - 1} \right| \right]. \quad (16)$$

It has a form similar to the one of the cross section for scattering on two Coulomb centres [1]. For equally charged monopoles $eg_1 = eg_2 = n/2$ we have

$$p_{1,2}^f = \pm n/R$$

and

$$\sigma = \frac{n^2 p_0^2}{2|p|^4} \cdot \left(1 + \left| \frac{1}{1 - p^2 \cdot R^2/n^2} \right| \right). \quad (17)$$

For monopole-antimonopole pair with $eg_1 = -eg_2 = n/2$ one obtains:

$$\sigma = \frac{nRp_0^2}{2|p|^3} \cdot \frac{1}{|1 - iRp/n|}, \quad p_1^f = 0, \quad p_2^f = -\frac{in}{R}. \quad (18)$$

In the limit of small $|p| \ll \frac{n}{R}$ one gets a scattering on magnetic dipole with moment $d = nR/e$.

3 Quantum conformal scattering

Let us now consider scattering on a system of electric and magnetic charges in the eikonal approximation. As mentioned above the theory becomes greatly

simplified for harmonic potentials [2]. In the eikonal approximation the scattering amplitude can be expressed as a double integral over the impact-parameter plane

$$f = \frac{p_0}{2\pi i} \int_{-\infty}^{\infty} dx \int_{-\infty e^{i\epsilon}}^{\infty e^{-i\epsilon}} dy \exp \left[-i(xp_x + yp_y) + \frac{i}{p_0} V(x, y) \right], \quad (19)$$

where

$$V(x, y) = \int_{-\infty}^{\infty} dz U(x, y, z) - \frac{ep_0}{c} \int_{-\infty}^{\infty} dz A_z(x, y, z) \quad (20)$$

is, as previously, a harmonic function. To pass to the classical limit one may assume the argument of the exponential in (19) to be large and evaluate the integral by the saddle point method.

The potential $V(x, y)$ is a harmonic function, and hence it can be represented as a real part of analytical function of the variables $b = x + iy$ (the complex impact parameter) and $b^* = x - iy$, where x and y are real:

$$V(x, y) = \left[V(b) + V^*(b^*) \right]. \quad (21)$$

It was shown in [2] that passing from variables x and y to b and b^* leads to factorization of the integrand and to the separation of variables b and b^* which enables us to express the result as product of contour integrals

$$f = \frac{p_0}{4\pi} \sum_{j=1}^{M+N} I_j(p) I_j(p^*), \quad (22)$$

where $I(p), I(p^*)$ are functions of the complex momentum $p = p_x + ip_y$ and of the conjugate momentum $p^* = p_x - ip_y$, respectively, and the sum goes over positions of charges (electric and magnetic):

$$I(p) I(p^*) = \int_{C_{b_j^*}} \exp \left[-\frac{1}{2} ib^* p + \frac{i}{p_0} V^*(b^*) \right] db^* \int_{C_{b_j}} \exp \left[-\frac{1}{2} ib p^* + \frac{i}{p_0} V(b) \right] db. \quad (23)$$

The cuts in the complex impact-parameter plane (b) go parallel to the real axis from $-\infty$ to each of the points b_j and b_j^* (positions of charges). The form of contours $C_{b_j^*}$ and C_{b_j} are shown in Fig.1.

Hence for the problem of small-angle scattering by a system of electric and magnetic charges the scattering amplitude is expressed as a product of two similar contour integrals, one of them depending on p and the other on p^* .

3.1 Special cases

Let us apply the technique of conformal scattering to some special cases. First we consider a single dyon with charges (q, g) located at $b_0 = x_0 + iy_0$ and a Dirac string characterized by the angle θ . The evaluation of contour integrals $I(p)$ and $I(p^*)$ given by equation (23) with

$$V(b) = \left(eq + i \frac{egp_0}{c} \right) \ln[(b - b_0) e^{i\theta}],$$

gives the eikonal amplitude for scattering on a dyon:

$$\begin{aligned} f(p) = & 2i \frac{p_0}{|p|^2} \left(\sqrt{\frac{p}{p^*}} \right)^{eg/c} \left(\frac{2p_0}{|p|} \right)^{2iq/p_0} \left(\frac{iq}{p_0} + \frac{eg}{c} \right) \times \\ & \times \exp(-i\text{Re}(b_0 p^*) - i\pi eg/c + 2ieg\theta) \frac{\Gamma\left(\frac{iq}{p_0} + \frac{eg}{c}\right)}{\Gamma\left(\frac{iq}{p_0} - \frac{eg}{c}\right)}. \end{aligned} \quad (24)$$

We see that the dependence on the direction of the Dirac string θ enters only as a constant phase and hence it does not affect any physical observables. It is easy to show that this simple dependence on the Dirac-strings directions is also pertinent for more general cases of many monopoles. In the case of a system containing M Dirac monopoles with charges g_i ($i = 1, \dots, M$) and “Dirac-strings angles” θ_i ($i = 1, \dots, M$) the dependence on the latter has a simple form of a constant phase:

$$e^{i \sum_{i=1}^M 2eg_i \theta_i / c}.$$

Using this fact one can set all $\theta_i = 0$ in all our calculations. The expression (24) for small-angle scattering amplitude on a dyon coincides with the result of rather

tedious calculations of ref [8]. The corresponding effective cross section on the dyon takes the following form, coinciding with the classical cross section (11)

$$\sigma = \frac{4}{|p|^4} \cdot \left\{ q^2 + \left(\frac{egp_0}{c} \right)^2 \right\}. \quad (25)$$

In the special case of a Dirac monopole $e \cdot g = 1/2$ placed at $b = 0$ we obtain the small-angle scattering amplitude:

$$f = -\frac{n}{\pi} \frac{p_0}{|p|^2} \left(\frac{p}{p^*} \right)^n \left(\frac{\tau^*}{\tau} \right)^{\frac{n}{2}} \exp \left[-\frac{i}{4} \text{Re} \left((b_2 + b_1) p^* \right) \right] \cdot \left[K_{\frac{1+n}{2}} \left(\frac{\tau^*}{2} \right) \cdot K_{\frac{1-n}{2}} \left(-\frac{\tau}{2} \right) - K_{\frac{1+n}{2}} \left(-\frac{\tau^*}{2} \right) \cdot K_{\frac{1-n}{2}} \left(\frac{\tau}{2} \right) \right]. \quad (26)$$

$$f(p) = \frac{p_0}{|p|^2} \sqrt{\frac{p}{p^*}} \left(\frac{2p_0}{|p|} \right)^{2iq/p_0} \left(1 + \frac{2iq}{p_0} \right) \frac{\Gamma \left(\frac{1}{2} + \frac{iq}{p_0} \right)}{\Gamma \left(\frac{1}{2} - \frac{iq}{p_0} \right)}. \quad (27)$$

Now one can relatively easy calculate the amplitude for scattering on two monopoles in the eikonal approximation. Let the complex numbers b_1 and b_2 be the projections of the magnetic charges g_1 and g_2 onto the impact-parameter plane. Introducing the notations:

$$\begin{aligned} \mathfrak{ae} &= \frac{e}{2}(g_2 - g_1), \\ \mu &= \frac{1}{2} - \frac{e}{2}(g_2 + g_1), \quad \mu' = 1 - \mu, \\ \tau &= \frac{ip}{2}(b_2 - b_1)^*, \end{aligned}$$

after simple calculations we obtain the following expressions for integrals (23) in the terms of the Whittaker function [6] $W_{\mathfrak{ae}, \mu}$:

$$\begin{aligned} I(p) &= 2\pi i (-1)^{-\mathfrak{ae}} e^{-\frac{1}{4}ip(b_1^* + b_2^*)} (b_1^* - b_2^*)^{2\mu} \tau^{-\mu - \frac{1}{2}} \frac{1}{\Gamma \left(\mathfrak{ae} - \mu + \frac{1}{2} \right)} W_{\mathfrak{ae}, \mu}(-\tau), \\ I(p^*) &= e^{-\frac{1}{4}ip^*(b_1 + b_2)} (b_1 - b_2)^{2\mu'} \tau^{*- \mu' - \frac{1}{2}} \Gamma \left(-\mathfrak{ae} + \mu' + \frac{1}{2} \right) W_{-\mathfrak{ae}, \mu'}(\tau^*). \end{aligned}$$

Implying Dirac's quantization conditions $eg_1 = n_1/2$, $eg_2 = n_2/2$, where $(n_1, n_2) \in \mathbf{Z}$, one gets the amplitude for scattering on two monopoles:

$$\begin{aligned}
f(p) = & -\frac{ip_0}{|p|^2} \left(-\frac{p^*}{p}\right)^{\frac{n_1+n_2}{2}} \left(\frac{\tau}{\tau^*}\right)^{\frac{n_1+n_2}{4}} \exp\left\{-\frac{i}{4} \operatorname{Re}((b_1 + b_2)p^*)\right\} \times \\
& \times \left\{(-1)^{n_1/2 - \frac{n_1+n_2}{4}} \cdot n_1 W_{\frac{n_1-n_2}{4}, \frac{1}{2} - \frac{n_2+n_1}{4}}(\tau^*) \cdot W_{\frac{n_2-n_1}{4}, \frac{1}{2} + \frac{n_2+n_1}{4}}(-\tau) + \right. \\
& \left. + (-1)^{n_2/2 + \frac{n_1+n_2}{4}} \cdot n_2 W_{-\frac{n_1-n_2}{4}, \frac{1}{2} - \frac{n_2+n_1}{4}}(-\tau^*) \cdot W_{-\frac{n_2-n_1}{4}, \frac{1}{2} + \frac{n_2+n_1}{4}}(\tau)\right\}. \quad (28)
\end{aligned}$$

The formula for the eikonal amplitude for the scattering by two identical monopoles $n_1 = n_2 = n$ simplifies, in this is case $\alpha = 0$ and hence we can use the representations for the Whittaker functions in terms of modified Bessel function $K_\mu(z)$:

$$W_{0,\mu}(z) = \sqrt{\frac{z}{\pi}} K_\mu\left(\frac{z}{2}\right).$$

Using this relation one obtains the scattering amplitude for small-angle scattering on two identical monopoles with charges $eg = n/2$:

$$\begin{aligned}
f = & \frac{n}{2\pi} \frac{p_0|b_2 - b_1|}{|p|} \left(-\frac{p^*}{p}\right)^n \left(\frac{\tau^*}{\tau}\right)^{\frac{n}{2}} \exp\left\{-\frac{i}{4} \operatorname{Re}((b_2 + b_1)p^*)\right\} \times \\
& \times \left[K_{\frac{1-n}{2}}\left(\frac{\tau^*}{2}\right) \cdot K_{\frac{1+n}{2}}\left(-\frac{\tau}{2}\right) - (-1)^n K_{\frac{1-n}{2}}\left(-\frac{\tau^*}{2}\right) \cdot K_{\frac{1+n}{2}}\left(\frac{\tau}{2}\right) \right]. \quad (29)
\end{aligned}$$

The amplitude for the case of small-angle scattering on M monopoles is expressed in terms of generalized hypergeometric functions.

4 Focal points

The small angle classical cross section for scattering of a charge particle on a system of point-like electric and magnetic charges diverges at particular values of the momentum transfer (focal points p_f) determined by eq. (2). In the quantum case this divergence is shadowed by quantum corrections, but in the semiclassical

limit the amplitude has clear maxima in a neighbourhood of focal points. The semiclassical amplitude in the neighbourhood of the focal points is not given by the steepest descent method for integral (19). For spherically symmetric scatterers the semiclassical amplitude in the neighbourhood of the focal points is given by the Airy function [9]. The more complicated case of two-centre Coulomb scatterer was considered in [2]. It was shown there that the corresponding amplitude in the neighbourhood of the focal points is described by canonical integral, which is called elliptic umbilica [7] in catastrophe theory.

We consider in details the approach to the semiclassical limit of the amplitudes in the neighbourhood of a focal point for two cases: scattering on two Coulomb centres and on two monopoles. Let us consider two systems: the first one with two equal electric charges q placed and $b = \pm R$, the second one is a system of two equal magnetic charges with $eg = n/2$ ($n \in \mathbf{Z}$). The values of the momentum transfer corresponding to focal points are given for the first system [2] by $p_f = \pm 2iq/p_0 R$ and for the second one by $p_f = \pm n/R$ (see section 2). The scattering becomes semiclassical if the following conditions are satisfied: $p_0 R \gg 1$ (incident particles wavelength much less than the size of the system) and $\theta_f = \left| \frac{p_f}{p_0} \right| \gg 1$ (classical scattering angle larger than the quantum mechanical uncertainty of the scattering angle). The above conditions imply for the first system that $q/p_0 \gg 1$ and $n \gg 1$ for the second one. In Fig.2 we show the appearance of maxima in cross sections corresponding to focal points for both systems with approaching to the classical limit at increasing values of q/p_0 and n for the first and the second system correspondingly.² The cross sections in the neighbourhood of the focal points are shown in Fig.3, one sees that for both systems the cross sections have clear maxima and have a similar behaviour corresponding to elliptic umbilica.

²For electric case we plot $|f|^2$ and for magnetic case we plot $|pf|^2$ to see focal more clearly.

5 CONCLUSION

The method of conformal scattering is generalized to the case of systems involving magnetic charges besides the electric ones. Calculations of the scattering amplitudes in the eikonal approximation are simplified considerably due to conformal properties of mapping of the impact parameter plane to the momentum transfer plane. This property enables us to introduce the notion of two dimensional complex analytical potential. The complex potential for a system involving magnetic charges corresponds to a system of Coulomb centres with complex charges having imaginary part proportional to the magnetic charge and the momentum of incident particle. Using these simplifications we obtained an explicit expression for the small-angle scattering amplitude on a system of two monopoles in terms of confluent hypergeometric functions for generic case and Bessel functions for two equal magnetic charges.


These explicit formulas enable us to study the semiclassical limit of the corresponding amplitudes in the neighbourhood of rainbow singularities (focal points). The behaviour of the cross-sections in the neighbourhood of these point singularities has a form of elliptic umbilica catastrophe.

6 ACKNOWLEDGMENT

This work has been supported by Deutscher Akademischer Austauschdienst (DAAD). The author is indebted to Prof. Yu. Demkov for numerous discussions of method of conformal scattering. The author is sincerely grateful to Prof. K. Goeke for warm hospitality.

References


- [1] Yu.N.Demkov, 1981 Sov. Phys. JETP **53(1)**, 63
- [2] D.I.Abramov, Yu.N.Demkov, and A.P.Scherbakov, 1981 Sov. Phys. JETP **53(4)**, 681
- [3] P.A.M. Dirac, 1931 Proc.Roy.Soc. (London) **A133**, 60; 1948 Phys.Rev. **74**, 817.
- [4] L.D.Landau, E.M.Lifshitz, 1974 Quantum Mechanics, M., Nauka,
- [5] I.S.Gradshteyn, I.M.Ryzhik, 1965 Table of Integrals, Series, and Products, 1064, New York: Academic Press
- [6] A.Erdelyi, 1953 Higher Transcendental Functions, vol.II, 116, McGraw-Hill
- [7] M.V.Berry, J.F.Nye, and F.J.Wright, 1979 Phil. Trans. Roy.Soc. **A291**, 453
- [8] G. Calucci, 1981 Lett. Nuovo Cim. **32**, 205
- [9] Roger G. Newton, 1953 Scattering theory of waves and particle, New-York:McGraw-Hill




M. Penttinen,

Small-angle scattering on a system of magnetic and electric charges

Fig.1



M. Penttinen,
 Small-angle scattering on a system of magnetic and electric charges,
Fig.2



M. Penttinen,

Small-angle scattering on a system of magnetic and electric charges,

Fig.3